

action of BK (10-1000 ng).

However, clotrimazole (1 μ M), an inhibitor of P450 reduced responses up to 80% while 7-ethoxyresorufin, another P450 inhibitor, was less effective (40% inhibition). 17-ODYA (2 μ M), an inhibitor of P450 fatty acid metabolism, also reduced responses to BK (up to 50%) suggesting a role of AA. None of the inhibitors affected responses to the reference vasodilator, nitroprusside (NP, 1000 ng). Vasodilator responses to BK, but not NP, were markedly reduced by 10mM TEA (85%) and procaine (80%) suggesting an effect mediated by increased K^+ conductance. Nifedipine (5 nM) almost abolished responses to BK and cromakalim (1-10 μ g) but did not affect those to NP. Inhibition of ATP-sensitive K^+ channels with glibenclamide (10 μ M) reduced responses to cromakalim but not those to BK. These results suggest that the coronary vasodilator response to BK is mediated by a P450-AA metabolite that stimulates a Ca^{2+} -activated K^+ channel.

References:

- 1 Oyekan AO, McGiff JC, Rosencrantz-Weiss P, Quilley J. Relaxant responses of rabbit aorta: influence of cytochrome P450 inhibitors. *J Pharmacol Exp Ther* 1994; 268: 262-269.
- 2 Fulton D, McGiff JC, Quilley J. Contribution of NO and cytochrome P450 to the vasodilator effect of bradykinin in the rat kidney. *Br J Pharmacol* 1992; 107: 722-725.
- 3 Oyekan AO, McGiff JC, Quilley J. Cytochrome P450-dependent vasodilation of rat kidney by arachidonic acid. *Am J Physiol* 1991; 261: H714-H719.

.....

25

CALPONIN PHOSPHORYLATION AND VASCULAR SMOOTH MUSCLE CONTRACTION

T. Tanaka, T. Mino, U. Yuasa, M. Naka

Department of Molecular and Cellular Pharmacology Mie University School of Medicine, Tsu, Mie 514, Japan

In response to many agonists, both inositol(1,4,5)-triphosphate (InsP3) and diacylglycerol (DAG) are formed by the hydrolysis of an inositol lipid precursor stored in the plasma membrane of smooth muscle. The InsP3 released into the cytoplasm mobilizes calcium from internal stores, whereas DAG stimulates protein kinase C. Calponin is a thin filament-associated smooth muscle protein that has been implicated to play a role in the regulation of smooth muscle [1]. Recently, we found that smooth muscle calponin is an excellent substrate for protein kinase C and the phosphorylation reduced the binding of calponin to F-actin and tropomyosin [2-4]. We have identified an important phosphorylation site in calponin by protein kinase C and demonstrated the calponin phosphorylation response following stimulation by endothelin-1 or phorbol 12,13-dibutyrate (PDBu) in 32P-labeled porcine coronary artery [5]. We found that Thr184 is the preferred site of phosphorylation and is functionally the most important of the sites phosphorylated by protein kinase C in smooth muscle calponin. We investigated the calponin phosphorylation during endothelin-1 or PDBu stimulation of intact strips of porcine coronary artery. Stimulation by endothelin-1 or PDBu resulted in a significant increase of 32P incorporation into the calponin in association with development of force. These results suggest that calponin phosphorylation plays a potential role in the regulation of smooth muscle contraction by endothelin-1 or PDBu.

References:

- 1 Takahashi K, Hiwada K, Kokuba T. Isolation and Characterization of a 34,000-dalton calmodulin- and F-actin-binding protein from chicken gizzard smooth muscle. *Biochem Biophys Res Commun* 1986; 141: 20-6.
- 2 Naka M, Kureishi Y, Muroga Y, Takahashi K, Ito M, Tanaka T. Modulation of smooth muscle calponin by protein kinase C and calmodulin. *Biochem Biophys Res Commun* 1990; 171: 933-937.
- 3 Winder SJ, Walsh MP. Smooth muscle calponin. Inhibition of actomyosin MgATPase and regulation by phosphorylation. *J Biol Chem* 1990; 265: 10148-55.
- 4 Nakamura F, Mino T, Yamamoto J, Naka M, Tanaka T. Identification of the regulatory site in smooth muscle calponin that is phosphorylated by protein kinase C. *J Biol Chem* 1993; 268: 6194-6201.
- 5 Ichikawa K, Ito M, Okubo S, Konishi T, Nakano T, Mino T, Nakamura F, Naka M, Tanaka T. Calponin phosphatase from smooth muscle: a possible role of type 1 protein phosphatase in smooth muscle relaxation. *Biochem Biophys Res Commun* 1993; 193: 827-833.

.....

26

INTERACTION OF NITROXIDERGIC NERVE WITH CHOLINERGIC AND NORADRENERGIC NERVES IN BLOOD VESSEL

N. Toda

Department of Pharmacology, Shiga University of Medical Sciences, Ohtsu 520-21, Japan

Nitroxicergic nerve in dog middle cerebral arteries originates from the pterygopalatine ganglion. VIPergic and cholinergic nerve cells and fibres are histochemically detected in the rat pterygopalatine ganglion which supplies nerve fibres to cerebral arteries. Nitroxicergic, cholinergic and VIPergic nerves are expected to be originated from the ganglion. Thus, we investigated functional the interrelationship between these nerves in isolated bovine basilar arteries. Transmural electrical stimulation (TES) relaxed the arteries. The response was not reduced in the VIP- and CGRP-tolerant arteries. Evidence for a mediation by NO of the response was obtained. The relaxation was reduced by acetylcholine or physostigmine and potentiated by atropine. VIP did not alter the response to TES. It is concluded that acetylcholine from cholinergic nerves appears to act on prejunctional muscarinic receptors and decrease the synthesis or release of NO in the nerve. VIP cannot be a transmitter nor a modulator of nitroxicergic nerve functions. In dog mesenteric arteries treated with α -adrenoceptor blocker, TES produced relaxation, which was proved to be mediated by NO. Treatment with EDRF or NO synthase inhibitors did not alter the release of norepinephrine by TES. These findings suggest that NO derived from the nerve does not interfere with the amine release from noradrenergic nerves.

References:

- 1 Toda N, Okamura T. Regulation by nitroxicergic nerve of arterial tone. *News Physiol Sci* 1992; 7: 148-152.
- 2 Toda N, Ayajiki K, Kimura H, Okamura T. Impairment by damage of the pterygopalatine in canine cerebral and retinal arteries. *Circ Res* 1993; 72: 206-213.
- 3 Ayajiki K, Okamura T, Toda N. Nitric oxide mediates, and acetylcholine modulates, neurally induced relaxation of bovine cerebral arteries. *Neuroscience* 1993; 54: 819-825.